

# Responsible and Effective Opioid Prescribing

## HOW TO RECEIVE CREDIT

- Read the enclosed course.
- Complete the questions at the end of the course.
- Return your completed Answer Sheet/Evaluation to NetCE by mail or fax, or complete online at [www.NetCE.com](http://www.NetCE.com). Your postmark or facsimile date will be used as your completion date.
- Receive your Certificate(s) of Completion by mail, fax, or email.

## Faculty Disclosure

Contributing faculty, Mark Rose, BS, MA, LP, has disclosed no relevant financial relationship with any product manufacturer or service provider mentioned.

## Division Planner

William E. Frey, DDS, MS, FICD

## Director of Development and Academic Affairs

Sarah Campbell

## Division Planner/Director Disclosure

The division planner and director have disclosed no relevant financial relationship with any product manufacturer or service provider mentioned.

## Audience

This course is designed for dental professionals who may alter prescribing practices or intervene to prevent drug diversion and inappropriate opioid use.

## Accreditations & Approvals

NetCE is an ADA CERP Recognized Provider.

ADA CERP is a service of the American Dental Association to assist dental professionals in identifying quality providers of continuing dental education. ADA CERP does not approve or endorse individual courses or instructors, nor does it imply acceptance of credit hours by boards of dentistry.

Concerns or complaints about a CE provider may be directed to the provider or to ADA CERP at [www.ada.org/cerp](http://www.ada.org/cerp).

NetCE

Nationally Approved PACE Program  
Provider for FAGD/MAGD credit.

Approval does not imply acceptance by  
any regulatory authority or AGD endorsement.  
10/1/2021 to 9/30/2027

Provider ID #217994.



## Faculty

**Mark Rose, BS, MA, LP**, is a licensed psychologist in the State of Minnesota with a private consulting practice and a medical research analyst with a biomedical communications firm. Earlier healthcare technology assessment work led to medical device and pharmaceutical sector experience in new product development involving cancer ablative devices and pain therapeutics. Along with substantial experience in addiction research, Mr. Rose has contributed to the authorship of numerous papers on CNS, oncology, and other medical disorders. He is the lead author of papers published in peer-reviewed addiction, psychiatry, and pain medicine journals and has written books on prescription opioids and alcoholism published by the Hazelden Foundation. He also serves as an Expert Advisor and Expert Witness to law firms that represent disability claimants or criminal defendants on cases related to chronic pain, psychiatric/substance use disorders, and acute pharmacologic/toxicologic effects. Mr. Rose is on the Board of Directors of the Minneapolis-based International Institute of Anti-Aging Medicine and is a member of several professional organizations.

NetCE is a Registered Provider with the Dental Board of California. Provider number RP3841. Completion of this course does not constitute authorization for the attendee to perform any services that he or she is not legally authorized to perform based on his or her permit type.

NetCE is approved as a provider of continuing education by the Florida Board of Dentistry, Provider #50-2405.

### **Designations of Credit**

NetCE designates this activity for 3 continuing education credits.

AGD Subject Code 340.

This course meets the Dental Board of California's requirements for 3 units of continuing education.

Dental Board of California course #03-3841-00322.

### **About the Sponsor**

The purpose of NetCE is to provide challenging curricula to assist healthcare professionals to raise their levels of expertise while fulfilling their continuing education requirements, thereby improving the quality of healthcare.

Our contributing faculty members have taken care to ensure that the information and recommendations are accurate and compatible with the standards generally accepted at the time of publication. The publisher disclaims any liability, loss or damage incurred as a consequence, directly or indirectly, of the use and application of any of the contents. Participants are cautioned about the potential risk of using limited knowledge when integrating new techniques into practice.

### **Disclosure Statement**

It is the policy of NetCE not to accept commercial support. Furthermore, commercial interests are prohibited from distributing or providing access to this activity to learners.

### **Course Objective**

The purpose of this course is to provide dental professionals who prescribe or distribute opioids with an appreciation for the complexities of opioid prescribing and the dual risks of litigation due to inadequate pain control and drug diversion or misuse in order to provide the best possible patient care and to prevent a growing social problem.

### **Learning Objectives**

Upon completion of this course, you should be able to:

1. Apply epidemiologic trends in opioid use and misuse to current practice so at-risk patient populations can be more easily identified, assessed, and treated.
2. Create comprehensive treatment plans for patients with pain that address patient needs as well as drug diversion prevention.
3. Evaluate behaviors that may indicate drug seeking or diverting as well as approaches for patients suspected of misusing opioids.
4. Identify state and federal laws governing the proper prescription and monitoring of controlled substances.
5. Describe the available treatment modalities for opioid use disorder.



Sections marked with this symbol include evidence-based practice recommendations. The level of evidence and/or strength of recommendation, as provided by the evidence-based source, are also included so you may determine the validity or relevance of the information. These sections may be used in conjunction with the course material for better application to your daily practice.

## INTRODUCTION

Pain is the leading reason for seeking medical care, and pain management is a large part of many healthcare professionals' practice. Opioid analgesics are approved by the U.S. Food and Drug Administration (FDA) for moderate and severe pain and are broadly accepted in acute pain, cancer pain, and end-of-life care, but are controversial in chronic noncancer pain. In response to the long-standing neglect of severe pain, indications for opioid analgesic prescribing were expanded in the 1990s, followed by inappropriate prescribing and increasing abuse, addiction, diversion, and overdose through the 2000s. In tandem with the continued under-treatment of pain, these practice patterns led to needless suffering from uncontrolled pain, opioid analgesic addiction, and overdose. Opioid analgesic prescribing and associated overdose peaked in 2011 with both now in multi-year decline.

Patients show substantial opioid response variations in analgesia and tolerability and may exhibit a range of psychologic, emotional, and behavioral responses that reflect inadequate pain control, an emerging opioid use problem, or both. Clinician delivery of best possible care to patients with pain requires appreciation of the complexities of opioid prescribing and the dual risks of inadequate pain control and inappropriate use, drug diversion, or overdose. A foundation for appropriate opioid prescribing is the understanding of factual data that clarify the prevalence, causality, and prevention of serious safety concerns with opioid prescribing.

## SCOPE OF THE PROBLEM

Inappropriate opioid analgesic prescribing for pain is defined as the non-prescribing, inadequate prescribing, excessive prescribing, or continued prescribing despite evidence of ineffectiveness of opioids [1]. Appropriate opioid prescribing is essential to achieve pain control; to minimize patient risk of abuse, addiction, and fatal toxicity; and to minimize societal harms from diversion.

The foundation of appropriate opioid prescribing is thorough patient assessment, treatment planning, and follow-up and monitoring. Essential for proper patient assessment and treatment planning is comprehension of the clinical concepts of opioid abuse and addiction, their behavioral manifestations in patients with pain, and how these potentially problematic behavioral responses to opioids both resemble and differ from physical dependence and pseudo-dependence. Prescriber knowledge deficit has been identified as a key obstacle to appropriate opioid prescribing and, along with gaps in policy, treatment, attitudes, and research, contributes to widespread inadequate treatment of pain [2].

The extent of opioid analgesic use in the United States in the 2000s was unprecedented in the country's history and unparalleled anywhere in the world. Before 1990, physicians in the United States were skeptical of prescribing opioids for chronic noncancer pain. In 2017, 20% of adults are prescribed an opioid such as oxycodone and hydrocodone for chronic pain, and sales of opioid analgesics totaled approximately \$7 billion in 2016 [10; 33].

Worldwide consumption of opioid analgesics has increased dramatically in the past few decades, with the United States driving a substantial proportion of this increase. For example, the 1990 global consumption of hydrocodone was 4 tons (3,628 kg), compared with the 2009 consumption of 39 tons (35,380 kg); 99% of this was consumed in the United States. Similarly, 3 tons (2,722 kg) of oxycodone were consumed globally in 1990, versus 77 tons (69,853 kg) in 2009, of which 62 tons (56,245 kg or 81%) were consumed in the United States [3]. With only 4.5% of the world's population, the United States annually consumes more than 80% of all opioid supplies, including [4]:

- 99% of all hydrocodone
- 80% of all oxycodone
- 58% of all methadone
- 54% of all hydromorphone
- 49% of all fentanyl
- 43% of all meperidine

This disproportionate rate of opioid consumption reflects sociocultural and economic factors and standards of clinical medicine.

Between 1992 and 2003, the U.S. population increased 14%, while persons abusing opioid analgesics increased 94% and first-time non-medical opioid analgesic users 12 to 17 years of age increased 542% [4]. It is interesting to note that while opioid prescribing has increased precipitously among adults in the United States, the rate remained low and steady for children between 1996 and 2012 [5]. A study using data from 2005 to 2015 showed opioid prescribing in 57 million visits from adolescents and young adults, representing a prescribing rate of nearly 15% in emergency departments and nearly 3% in outpatient clinical settings [36]. During the course of the study, emergency department prescribing decreased slightly while outpatient clinical setting prescribing remained the same [36]. To assist in monitoring the public health problem associated with prescribed opioids, numerous governmental, non-profit, and private sector agencies and organizations are involved in collecting, reporting, and analyzing data on the abuse, addiction, fatal overdose, and treatment admissions related to opioid analgesics.

Before it was halted in 2011, the Drug Abuse Warning Network (DAWN) provided estimates of the health consequences of nonmedical use of individual drugs, including opioid medications [6]. DAWN indicates that opioid abuse is a growing problem in the United States. In 2005 and 2011, hydrocodone and its combinations accounted for 51,225 and 97,183 emergency department visits, respectively. Oxycodone and its combinations resulted in 42,810 visits to the emergency department in 2005; this number increased to 175,229 visits in 2011 [7; 8]. Visits for nonmedical use of all opioids increased from 217,594 to 420,040 during the six-year period. In 2016–2017, there were 127,101 nonmedical opioid emergency department visits [39]. While this number is an improvement from previous years, nonmedical use accounts for 47.6% of all emergency department visits related to opioids [39].

---

## PAIN MANAGEMENT APPROACHES

---

Healthcare professionals should know the best clinical practices in opioid prescribing, including the associated risks of opioids, approaches to the assessment of pain and function, and pain management modalities. Pharmacologic and nonpharmacologic approaches should be used on the basis of current knowledge in the evidence base or best clinical practices. Patients with moderate-to-severe chronic pain who have been assessed and treated, over a period of time, with non-opioid therapy or nonpharmacologic pain therapy without adequate pain relief, are considered to be candidates for a trial of opioid therapy [9; 10]. Initial treatment should always be considered individually determined and as a trial of therapy, not a definitive course of treatment [11].

In 2016, the CDC issued updated guidance on the prescription of opioids for chronic pain [10]. The guideline addresses when to initiate or continue opioids for chronic pain; opioid selection, dosage, duration, follow-up, and discontinuation; and assessing risk and addressing harms of opioid use. In addition, the CDC further updated guidance against the misapplication of this guideline in 2019, noting that some policies and practices attributed to the guideline were inconsistent with the recommendations [40].

### ACUTE PAIN

Long-term opioid use often begins with treatment of acute pain. When opioids are used for acute pain, clinicians should prescribe the lowest effective dose of immediate-release opioids in a quantity no greater than that needed for the expected duration of severe pain. In most cases, three days or less will be sufficient; more than seven days will rarely be needed [10]. However, it is important to note that this guideline is based on emergency department prescribing guidelines for non-traumatic non-surgical pain [12]. It may be necessary to prescribe for longer periods in patients with acute severe pain.

With postoperative, acute, or intermittent pain, analgesia often requires frequent titration, and the two- to four-hour analgesic duration with short-acting hydrocodone, morphine, and oxycodone is more effective than extended-release formulations. Short-acting opioids are also recommended in patients who are medically unstable or with highly variable pain intensity [13; 14; 15].

## CHRONIC PAIN

Nonpharmacologic therapy and non-opioid pharmacologic therapy are the preferred first-line therapies for chronic pain. Several nonpharmacologic approaches are therapeutic complements to pain-relieving medication, lessening the need for higher doses and perhaps minimizing side effects. These interventions can help decrease pain or distress that may be contributing to the pain sensation. Approaches include palliative radiotherapy, complementary/alternative methods, manipulative and body-based methods, and cognitive/behavioral techniques. The choice of a specific nonpharmacologic intervention is based on the patient's preference, which, in turn, is usually based on a successful experience in the past.

Implantable intrathecal opioid infusion and/or spinal cord stimulation may be options for severe, intractable pain. Both options require that devices or ports be implanted, with associated risks. With intrathecal opioid infusion, the ability to deliver the drug directly into the spine provides pain relief with significantly smaller opioid doses, which can help to minimize side effects (e.g., drowsiness, dizziness, dry mouth, nausea, vomiting, and constipation) that can accompany systemic pain medications that might be delivered orally, transdermally, or through an IV [43]. However, use of opioid infusion has traditionally been limited to cancer pain. With spinal cord stimulation therapy, the most challenging aspect is patient selection. In order for patients to be considered for spinal cord stimulation, other options should have been ineffective or be contraindicated. Spinal cord stimulation is indicated for severe neuropathic pain persisting at least six months [91].

If opioids are used, they should be combined with nonpharmacologic therapy and non-opioid pharmacologic therapy, as appropriate. Clinicians should consider opioid therapy only if expected benefits for pain and function are anticipated to outweigh risks to the patient [10].

Opioid therapy for chronic pain should be presented as a trial for a pre-defined period (e.g.,  $\leq 30$  days). The goals of treatment should be established with all patients prior to the initiation of opioid therapy, including reasonable improvements in pain, function, depression, anxiety, and avoidance of unnecessary or excessive medication use [1; 10]. The treatment plan should describe therapy selection, measures of progress, and other diagnostic evaluations, consultations, referrals, and therapies.

In patients who are opioid-naïve, start at the lowest possible dose and titrate to effect. Dosages for patients who are opioid-tolerant should always be individualized and titrated by efficacy and tolerability [1; 10]. When starting opioid therapy for chronic pain, clinicians should prescribe short-acting instead of extended-release/long-acting opioid formulations [10].

The need for frequent progress and benefit/risk assessments during the trial should be included in patient education. Patients should also have full knowledge of the warning signs and symptoms of respiratory depression. Prescribers should carefully reassess evidence of benefits and risks when increasing the dosage to  $\geq 50$  mg morphine equivalent dose (MED) per day. Decisions to titrate dose to  $\geq 90$  mg MED/day should be avoided or carefully justified [10; 40].

Prescribers should be knowledgeable of federal and state opioid prescribing regulations. Issues of equianalgesic dosing, close patient monitoring during all dose changes, and cross-tolerance with opioid conversion should be considered. If necessary, treatment may be augmented, with preference for nonopioids and immediate-release opioids over long-acting/extended-release opioids. Taper opioid dose when no longer needed [16].

## PALLIATIVE CARE AND PAIN AT THE END OF LIFE

Unrelieved pain is the greatest fear among people with a life-limiting disease, and the need for an increased understanding of effective pain management is well-documented [27]. Although experts have noted that 75% to 90% of end-of-life pain can be managed effectively, rates of pain are high, even among people receiving palliative care [27; 30; 34; 35].

The inadequate management of pain is the result of several factors related to both patients and clinicians. In a survey of oncologists, patient reluctance to take opioids or to report pain were two of the most important barriers to effective pain relief [37]. This reluctance is related to a variety of attitudes and beliefs [27; 37]:

- Fear of addiction to opioids
- Worry that if pain is treated early, there will be no options for treatment of future pain
- Anxiety about unpleasant side effects from pain medications
- Fear that increasing pain means that the disease is getting worse
- Desire to be a “good” patient
- Concern about the high cost of medications

Education and open communication are the keys to overcoming these barriers. Every member of the healthcare team should reinforce accurate information about pain management with patients and families. The clinician should initiate conversations about pain management, especially regarding the use of opioids, as few patients will raise the issue themselves or even express their concerns unless they are specifically asked [38]. It is important to acknowledge patients' fears individually and provide information to help them differentiate fact from fiction. For example, when discussing opioids with a patient who fears addiction, the clinician should explain that the risk of addiction is low [27]. It is also helpful to note the difference between addiction and physical dependence.

There are several other ways clinicians can allay patients' fears about pain medication:

- Assure patients that the availability of pain relievers cannot be exhausted; there will always be medications if pain becomes more severe.
- Acknowledge that side effects may occur but emphasize that they can be managed promptly and safely and that some side effects will abate over time.
- Explain that pain and severity of disease are not necessarily related.

Encouraging patients to be honest about pain and other symptoms is also vital. Clinicians should ensure that patients understand that pain is multidimensional and emphasize the importance of talking to a member of the healthcare team about possible causes of pain, such as emotional or spiritual distress. The healthcare team and patient should explore psychosocial and cultural factors that may affect self-reporting of pain, such as concern about the cost of medication.

Clinicians' attitudes, beliefs, and experiences also influence pain management, with addiction, tolerance, side effects, and regulations being the most important concerns [27; 34; 37; 42]. A lack of appropriate education and training in the assessment and management of pain has been noted to be a substantial contributor to ineffective pain management [37; 42]. As a result, many clinicians, especially primary care physicians, do not feel confident about their ability to manage pain in their patients [37; 42].

Clinicians require a clear understanding of available medications to relieve pain, including appropriate dosing, safety profiles, and side effects. If necessary, clinicians should consult with pain specialists to develop an effective approach.

Strong opioids are used for severe pain at the end of life [30; 34]. Morphine, buprenorphine, oxycodone, hydromorphone, fentanyl, and methadone are the most widely used in the United States [45]. Unlike nonopioids, opioids do not have a ceiling effect, and the dose can be titrated until pain is relieved or side effects become unmanageable. For patients who are opioid-naïve or who have been receiving low doses of a weak opioid, the initial dose should be low, and, if pain persists, the dose may be titrated up daily until pain is controlled.

More than one route of opioid administration will be needed by many patients during end-of-life care, but in general, opioids should be given orally, as this route is the most convenient and least expensive. The transdermal route is preferred to the parenteral route, although dosing with a transdermal patch is less flexible and so may not be appropriate for patients with unstable pain [34]. Intramuscular injections should be avoided because injections are painful, drug absorption is unreliable, and the time to peak concentration is long [34].

---

## CREATING A TREATMENT PLAN AND ASSESSMENT OF ADDICTION RISK

---

Information obtained by patient history, physical examination, and interview, from family members, a spouse, or state prescription drug monitoring program (PDMP), and from the use of screening and assessment tools can help the clinician to stratify the patient according to level of risk for developing problematic opioid behavioral responses (**Table 1**) [17; 28]. Low-risk patients receive the standard level of monitoring, vigilance, and care. Moderate-risk patients should be considered for an additional level of monitoring and provider contact, and high-risk patients are likely to require intensive and structured monitoring and follow-up contact, additional consultation with psychiatric and addiction medicine specialists, and limited supplies of short-acting opioid formulations [10; 26].

Before deciding to prescribe an opioid analgesic, clinicians should perform and document a detailed patient assessment that includes [1]:

- Pain indications for opioid therapy
- Nature and intensity of pain
- Past and current pain treatments and patient response
- Comorbid conditions
- Pain impact on physical and psychologic function
- Social support, housing, and employment
- Home environment (i.e., stressful or supportive)
- Pain impact on sleep, mood, work, relationships, leisure, and substance use
- Patient history of physical, emotional, or sexual abuse

If substance abuse is active, in remission, or in the patient's history, consult an addiction specialist before starting opioids [1]. In active substance abuse, do not prescribe opioids until the patient is engaged in treatment/recovery program or other arrangement made, such as addiction professional co-management and additional monitoring. When considering an opioid analgesic (particularly those that are extended-release or long-acting), one must always weigh the benefits against the risks of overdose, abuse, addiction, physical dependence and tolerance, adverse drug interactions, and accidental exposure by children [10; 16].

Screening and assessment tools can help guide patient stratification according to risk level and inform the appropriate degree of structure and monitoring in the treatment plan. It should be noted that despite widespread endorsement of screening tools used to help determine patient risk level, most tools have not been extensively evaluated, validated, or compared to each other, and evidence of their reliability is poor [17; 28].

RISK STRATIFICATION FOR PATIENTS PRESCRIBED OPIOIDS	
<b>Low Risk</b>	
Definable physical pathology with objective signs and reliable symptoms Clinical correlation with diagnostic testing, including MRI, physical examination, and interventional diagnostic techniques With or without mild psychologic comorbidity With or without minor medical comorbidity No or well-defined and controlled personal or family history of alcoholism or substance abuse Age 45 years or older High levels of pain acceptance and active coping strategies High motivation and willingness to participate in multimodal therapy and attempting to function at normal levels	
<b>Medium Risk</b>	
Significant pain problems with objective signs and symptoms confirmed by radiologic evaluation, physical examination, or diagnostic interventions Moderate psychologic problems, well controlled by therapy Moderate coexisting medical disorders that are well controlled by medical therapy and are not affected by chronic opioid therapy (e.g., central sleep apnea) Develops mild tolerance but not hyperalgesia without physical dependence or addiction History of personal or family history of alcoholism or substance abuse Pain involving more than three regions of the body Defined pathology with moderate levels of pain acceptance and coping strategies Willing to participate in multimodal therapy, attempting to function in normal daily life	
<b>High Risk</b>	
Widespread pain without objective signs and symptoms Pain involving more than three regions of the body Aberrant drug-related behavior History of alcoholism or drug misuse, abuse, addiction, diversion, dependency, tolerance, or hyperalgesia Major psychologic disorders Age younger than 45 years HIV-related pain High levels of pain exacerbation and low levels of coping strategies Unwilling to participate in multimodal therapy, not functioning close to a near normal lifestyle	
HIV = human immunodeficiency syndrome, MRI = magnetic resonance imaging.	
Source: [17; 28]	
Table 1	

 EVIDENCE-BASED PRACTICE RECOMMENDATION

Despite limited evidence for reliability and accuracy, screening for opioid use is recommended by the American Society of Interventional Pain Physicians, as it will identify opioid abusers and reduce opioid abuse.  
[\(<https://painphysicianjournal.com/2012/july/2012;%2015;S67-S116.pdf>. Last accessed April 28, 2021.\)](https://painphysicianjournal.com/2012/july/2012;%2015;S67-S116.pdf)

**Level of Evidence:** Limited (Evidence is insufficient to assess effects on health outcomes because of limited number or power of studies, large and unexplained inconsistency between higher-quality trials, important flaws in trial design or conduct, gaps in the chain of evidence, or lack of information on important health outcomes.)

## RISK ASSESSMENT TOOLS

### Opioid Risk Tool (ORT)

The Opioid Risk Tool (ORT) is a five-item, patient-administered assessment to help predict aberrant drug-related behavior. The ORT is also used to establish patient risk level through categorization into low, medium, or high levels of risk for aberrant drug-related behaviors based on responses to questions of previous alcohol/drug abuse, psychologic disorders, and other risk factors [18].

## Screener and Opioid Assessment for Patients with Pain-Revised (SOAPP-R)

The Screener and Opioid Assessment for Patients with Pain-Revised (SOAPP-R) is a patient-administered, 24-item screen with questions addressing history of alcohol/substance use, psychologic status, mood, cravings, and stress. Like the ORT, the SOAPP-R helps assess risk level of aberrant drug-related behaviors and the appropriate extent of monitoring [18; 19].

## Screening Instrument or Substance Abuse Potential (SISAP)

The Screening Instrument or Substance Abuse Potential (SISAP) tool is a self-administered, five-item questionnaire addressing history developed used to predict the risk of opioid misuse. The SISAP is used to identify patients with a history of alcohol/substance abuse and improve pain management by facilitating focus on the appropriate use of opioid analgesics and therapeutic outcomes in the majority of patients who are not at risk of opioid abuse, while carefully monitoring those who may be at greater risk [18].

## CAGE and CAGE-AID

The original CAGE (Cut down, Annoyed, Guilty, and Eye-opener) Questionnaire consisted of four questions designed to help clinicians determine the likelihood that a patient was misusing or abusing alcohol. These same four questions were modified to create the CAGE-AID (adapted to include drugs), revised to assess the likelihood of current substance abuse [20].

## Diagnosis, Intractability, Risk, and Efficacy (DIRE) Score

The Diagnosis, Intractability, Risk, and Efficacy (DIRE) risk assessment score is a clinician-rated questionnaire that is used to predict patient compliance with long-term opioid therapy [18; 21]. Patients scoring lower on the DIRE tool are poor candidates for long-term opioid analgesia.

## INFORMED CONSENT AND TREATMENT AGREEMENTS

The initial opioid prescription is preceded by a written informed consent or “treatment agreement” [1]. This agreement should address potential side effects, tolerance and/or physical dependence, drug interactions, motor skill impairment, limited evidence of long-term benefit, misuse, dependence, addiction, and overdose. Informed consent documents should include information regarding the risk/benefit profile for the drug(s) being prescribed. The prescribing policies should be clearly delineated, including the number/frequency of refills, early refills, and procedures for lost or stolen medications.

The treatment agreement also outlines joint physician and patient responsibilities. The patient agrees to using medications safely, refraining from “doctor shopping,” and consenting to routine urine drug testing (UDT). The prescriber’s responsibility is to address unforeseen problems and prescribe scheduled refills. Reasons for opioid therapy change or discontinuation should be listed. Agreements can also include sections related to follow-up visits, monitoring, and safe storage and disposal of unused drugs.

## PERIODIC REVIEW AND MONITORING

When implementing a chronic pain treatment plan that involves the use of opioids, the patient should be frequently reassessed for changes in pain origin, health, and function [1]. This can include input from family members and/or the state PDMP. During the initiation phase and during any changes to the dosage or agent used, patient contact should be increased. At every visit, chronic opioid response may be monitored according to the “5 A’s” [1; 23]:

- Analgesia
- Activities of daily living
- Adverse or side effects
- Aberrant drug-related behaviors
- Affect (i.e., patient mood)

Signs and symptoms that, if present, may suggest a problematic response to the opioid and interference with the goal of functional improvement include [24; 29]:

- Excessive sleeping or days and nights turned around
- Diminished appetite
- Short attention span or inability to concentrate
- Mood volatility, especially irritability
- Lack of involvement with others
- Impaired functioning due to drug effects
- Use of the opioid to regress instead of re-engaging in life
- Lack of attention to hygiene and appearance

The decision to continue, change, or terminate opioid therapy is based on progress toward treatment objectives and absence of adverse effects and risks of overdose or diversion [1]. Satisfactory therapy is indicated by improvements in pain, function, and quality of life. Brief assessment tools to assess pain and function may be useful, as may UDTs. Treatment plans may include periodic pill counts to confirm adherence and minimize diversion.

### Involvement of Family

Family members of the patient can provide the clinician with valuable information that better informs decision making regarding continuing opioid therapy. Family members can observe whether a patient is losing control of his or her life or becoming less functional or more depressed during the course of opioid therapy. They can also provide input regarding positive or negative changes in patient function, attitude, and level of comfort. The following questions can be asked of family members or a spouse to help clarify whether the patient's response to opioid therapy is favorable or unfavorable [24; 29]:

- Is the person's day centered around taking the opioid medication? Response can help clarify long-term risks and benefits of the medication and identify other treatment options.
- Does the person take pain medication only on occasion, perhaps three or four times per week? If yes, the likelihood of addiction is low.
- Have there been any other substance (alcohol or drug) abuse problems in the person's life? An affirmative response should be taken into consideration when prescribing.
- Does the person in pain spend most of the day resting, avoiding activity, or feeling depressed? If so, this suggests the pain medication is failing to promote rehabilitation. Daily activity is essential, and the patient may be considered for enrollment in a graduated exercise program
- Is the person in pain able to function (e.g., work, do household chores, play) with pain medication in a way that is clearly better than without? If yes, this suggests the pain medication is contributing to wellness.

### Assessment Tools

#### VIGIL

VIGIL is the acronym for a five-step risk management strategy designed to empower clinicians to appropriately prescribe opioids for pain by reducing regulatory concerns and to give pharmacists a framework for resolving ambiguous opioid analgesic prescriptions in a manner that preserves legitimate patient need while potentially deterring diverters. The components of VIGIL are:

- Verification: Is this a responsible opioid user?
- Identification: Is the identity of this patient verifiable?
- Generalization: Do we agree on mutual responsibilities and expectations?

PATIENT RISK LEVEL AND FREQUENCY OF MONITORING			
Monitoring Tool	Patient Risk Level		
	Low	Medium	High
Urine drug test	Every 1 to 2 years	Every 6 to 12 months	Every 3 to 6 months
State prescription drug monitoring program	Twice per year	Three times per year	Four times per year

Source: [47] Table 2

- Interpretation: Do I feel comfortable allowing this person to have controlled substances?
- Legalization: Am I acting legally and responsibly?

The foundation of VIGIL is a collaborative physician/pharmacist relationship [25].

#### **Current Opioid Misuse Measure (COMM)**

The Current Opioid Misuse Measure (COMM) is a 17-item patient self-report assessment designed to help clinicians identify misuse or abuse in patients being treated for chronic pain. Unlike the ORT and the SOAPP-R, the COMM identifies aberrant behaviors associated with opioid misuse in patients already receiving long-term opioid therapy [26]. Sample questions include: In the past 30 days, how often have you had to take more of your medication than prescribed? In the past 30 days, how much of your time was spent thinking about opioid medications (e.g., having enough, taking them, dosing schedule)?

#### **Pain Assessment and Documentation Tool (PADT)**

Guidelines by the CDC, the Federation of State Medical Boards (FSMB), and the Joint Commission stress the importance of documentation from both a healthcare quality and medicolegal perspective. Research has found widespread deficits in chart notes and progress documentation with patients with chronic pain receiving opioid ther-

apy, and the Pain Assessment and Documentation Tool (PADT) was designed to address these shortcomings [46]. The PADT is a clinician-directed interview, with most sections (e.g., analgesia, activities of daily living, adverse events) consisting of questions asked of the patient. However, the potential aberrant drug-related behavior section must be completed by the physician based on his or her observations of the patient.

#### **The Brief Intervention Tool**

The Brief Intervention Tool is a 26-item, “yes-no,” patient-administered questionnaire used to identify early signs of opioid abuse or addiction. The items assess the extent of problems related to drug use in several areas, including drug use-related functional impairment [22].

#### **Urine Drug Tests**

UDTs may be used to monitor adherence to the prescribed treatment plan and to detect unsanctioned drug use. They should be used more often in patients receiving addiction therapy, but clinical judgment is the ultimate guide to testing frequency (**Table 2**) [47]. The CDC recommends clinicians should use UDT before starting opioid therapy and consider UDT at least annually to assess for prescribed medications as well as other controlled prescription drugs and illicit drugs [10]. However, this recommendation was based on low-quality evidence that indicates little confidence in the effect estimate.

Initially, testing involves the use of class-specific immunoassay drug panels [1]. If necessary, this may be followed with gas chromatography/mass spectrometry for specific drug or metabolite detection. It is important that testing identifies the specific drug rather than the drug class, and the prescribed opioid should be included in the screen. Any abnormalities should be confirmed with a laboratory toxicologist or clinical pathologist. Immunoassay may be used point-of-care for “on-the-spot” therapy changes, but the high error rate prevents its use in major clinical decisions except with liquid chromatography coupled to tandem mass spectrometry confirmation.

Urine test results suggesting opioid misuse should be discussed with the patient using a positive, supportive approach. The test results and the patient discussion should be documented.

## **CONCURRENT USE OF BENZODIAZEPINES**

In 2019, 16% of persons who died of an opioid overdose also tested positive for benzodiazepines, a class of sedative medication commonly prescribed for anxiety, insomnia, panic attack, and muscle spasm [44]. Benzodiazepines work by raising the level of the neurotransmitter gamma-aminobutyric acid (GABA) in the brain. Common formulations include diazepam, alprazolam, and clonazepam. Combining benzodiazepines with opioids is unsafe because both classes of drug cause central nervous system depression and sedation and can decrease respiratory drive—the usual cause of overdose fatality. Both classes have the potential for drug dependence and addiction.

The CDC recommends that healthcare providers avoid prescribing benzodiazepines concurrently with opioids whenever possible [10]. If a benzodiazepine is to be discontinued, the clinician should taper the medication gradually, because abrupt withdrawal can lead to rebound anxiety and complications such as hallucinations, seizures, delirium tremens, and, in rare instances, death. A commonly used tapering schedule is a reduction of the benzodiazepine dose by 25% every one to two weeks [10].

## **CONSULTATION AND REFERRAL**

It is important to seek consultation or patient referral when input or care from a pain, psychiatry, addiction, or mental health specialist is necessary. Clinicians who prescribe opioids should become familiar with opioid addiction treatment options (including licensed opioid treatment programs for methadone and office-based opioid treatment for buprenorphine) if referral is needed [1].

Ideally, providers should be able to refer patients with active substance abuse who require pain treatment to an addiction professional or specialized program. In reality, these specialized resources are scarce or non-existent in many areas [1]. Therefore, each provider will need to decide whether the risks of continuing opioid treatment while a patient is using illicit drugs outweigh the benefits to the patient in terms of pain control and improved function [48].

## **MEDICAL RECORDS**

As noted, documentation is a necessary aspect of all patient care, but it is of particular importance when opioid prescribing is involved. All clinicians should maintain accurate, complete, and up-to-date medical records, including all written or telephoned prescription orders for opioid analgesics and other controlled substances, all written instructions to the patient for medication use, and the name, telephone number, and address of the patient’s pharmacy [1]. Good medical records demonstrate that a service was provided to the patient and that the service was medically necessary. Regardless of the treatment outcome, thorough medical records protect the prescriber.

## **PATIENT EDUCATION ON THE USE AND DISPOSAL OF OPIOIDS**

Patients and caregivers should be counseled regarding the safe use and disposal of opioids. As part of its mandatory Risk Evaluation and Mitigation Strategy (REMS) for extended-release/long-acting opioids, the U.S. Food and Drug Administration (FDA) has developed a patient counseling document with information on the patient’s specific

medications, instructions for emergency situations and incomplete pain control, and warnings not to share medications or take them unprescribed [16]. A copy of this form may be accessed online at <https://www.fda.gov/media/114694/download>.

When prescribing opioids, clinicians should provide patients with the following information [16]:

- Product-specific information
- Taking the opioid as prescribed
- Importance of dosing regimen adherence, managing missed doses, and prescriber contact if pain is not controlled
- Warning and rationale to never break or chew/crush tablets or cut or tear patches prior to use
- Warning and rationale to avoid other central nervous system depressants, such as sedative-hypnotics, anxiolytics, alcohol, or illicit drugs
- Warning not to abruptly halt or reduce the opioid without physician oversight of safe tapering when discontinuing
- The potential of serious side effects or death
- Risk factors, signs, and symptoms of overdose and opioid-induced respiratory depression, gastrointestinal obstruction, and allergic reactions
- The risks of falls, using heavy machinery, and driving
- Warning and rationale to never share an opioid analgesic
- Rationale for secure opioid storage
- Warning to protect opioids from theft
- Instructions for disposal of unneeded opioids, based on product-specific disposal information

There are no universal recommendations for the proper disposal of unused opioids, and patients are rarely advised of what to do with unused or expired medications [49]. According to the FDA, most medications that are no longer necessary or have

expired should be removed from their containers, mixed with undesirable substances (e.g., cat litter, used coffee grounds), and put into an impermeable, nondescript container (e.g., disposable container with a lid or a sealed bag) before throwing in the trash [50]. Any personal information should be obscured or destroyed. The FDA recommends that certain medications, including oxycodone/acetaminophen (Percocet), oxycodone (OxyContin tablets), and transdermal fentanyl (Duragesic Transdermal System), be flushed down the toilet instead of thrown in the trash [31; 50]. The FDA provides a free toolkit of materials (e.g., social media images, fact sheets, posters) to raise awareness of the serious dangers of keeping unused opioid pain medicines in the home and with information about safe disposal of these medicines. The Remove the Risk Outreach toolkit is updated regularly and can be found at <https://www.fda.gov/drugs/ensuring-safe-use-medicine/safe-opioid-disposal-remove-risk-outreach-toolkit> [31]. Patients should be advised to flush prescription drugs down the toilet only if the label or accompanying patient information specifically instructs doing so.

The American College of Preventive Medicine has established best practices to avoid diversion of unused drugs and educate patients regarding drug disposal [49]:

- Consider writing prescriptions in smaller amounts.
- Educate patients about safe storing and disposal practices.
- Give drug-specific information to patients about the temperature at which they should store their medications. Generally, the bathroom is not the best storage place. It is damp and moist, potentially resulting in potency decrements, and accessible to many people, including children and teens, resulting in potential theft or safety issues.
- Ask patients not to advertise that they are taking these types of medications and to keep their medications secure.

- Refer patients to community “take back” services overseen by law enforcement that collect controlled substances, seal them in plastic bags, and store them in a secure location until they can be incinerated. Contact your state law enforcement agency or visit <https://www.dea.gov> to determine if a program is available in your area.

## DISCONTINUING OPIOID THERAPY

The decision to continue or end opioid prescribing should be based on a physician-patient discussion of the anticipated benefits and risks. An opioid should be discontinued with resolution of the pain condition, intolerable side effects, inadequate analgesia, lack of improvement in quality of life despite dose titration, deteriorating function, or significant aberrant medication use [1; 10].

Clinicians should provide patients physically dependent on opioids with a safely structured tapering protocol. Withdrawal is managed by the prescribing physician or referral to an addiction specialist. Patients should be reassured that opioid discontinuation is not the end of treatment; continuation of pain management will be undertaken with other modalities through direct care or referral.

As a side note, cannabis use by patients with chronic pain receiving opioid therapy has traditionally been viewed as a treatment agreement violation that is grounds for termination of opioid therapy. However, some now argue against cannabis use as a rationale for termination or substantial treatment and monitoring changes, especially considering the increasing legalization of medical use at the state level [48].

## CONSIDERATIONS FOR NON-ENGLISH-PROFICIENT PATIENTS

For patients who are not proficient in English, it is important that information regarding the risks associated with the use of opioids and available resources be provided in their native language,

if possible. When there is an obvious disconnect in the communication process between the practitioner and patient due to the patient’s lack of proficiency in the English language, an interpreter is required. Interpreters can be a valuable resource to help bridge the communication and cultural gap between patients and practitioners. Interpreters are more than passive agents who translate and transmit information back and forth from party to party. When they are enlisted and treated as part of the interdisciplinary clinical team, they serve as cultural brokers who ultimately enhance the clinical encounter. In any case in which information regarding treatment options and medication/treatment measures are being provided, the use of an interpreter should be considered. Print materials are also available in many languages, and these should be offered whenever necessary.

---

## IDENTIFICATION OF DRUG DIVERSION/SEEKING BEHAVIORS

Research has more closely defined the location of prescribed opioid diversion into illicit use in the supply chain from the manufacturer to the distributor, retailer, and the end user (the pain patient). This information carries with it substantial public policy and regulatory implications. The 2019 National Survey on Drug Use and Health asked non-medical users of prescription opioids how they obtained their most recently used drugs [51]. Among persons 12 years of age or older, 38.6% obtained their prescription opioids from a friend or relative for free, 34.7% got them through a prescription from one doctor (vs. 17.3% in 2009–2010), 9.5% bought them from a friend or relative, and 3.2% took them from a friend or relative without asking [51]. Less frequent sources included a drug dealer or other stranger (6.5%); multiple doctors (2.0%); and theft from a doctor’s office, clinic, hospital, or pharmacy (0.9%) (vs. 0.2% in 2009–2010) [51].

As discussed, UDTs can give insight into patients who are misusing opioids. A random sample of UDT results from 800 patients treated for pain at a Veterans Affairs facility found that 25.2% were negative for the prescribed opioid while 19.5% were positive for an illicit drug/unreported opioid [52]. Negative UDT results for the prescribed opioid do not necessarily indicate diversion, but may indicate the patient halted his/her use due to side effects, lack of efficacy, or pain remission. The concern arises over the increasingly stringent climate surrounding clinical decision-making regarding aberrant UDT results and that a negative result for the prescribed opioid or a positive UDT may serve as the pretense to terminate a patient rather than guide him/her into addiction treatment or an alternative pain management program [53].

In addition to aberrant urine screens, there are certain behaviors that are suggestive of an emerging opioid use disorder. The most suggestive behaviors are [48; 54; 55]:

- Selling medications
- Prescription forgery or alteration
- Injecting medications meant for oral use
- Obtaining medications from nonmedical sources
- Resisting medication change despite worsening function or significant negative effects
- Loss of control over alcohol use
- Using illegal drugs or non-prescribed controlled substances
- Recurrent episodes of:
  - Prescription loss or theft
  - Obtaining opioids from other providers in violation of a treatment agreement
  - Unsanctioned dose escalation
  - Running out of medication and requesting early refills

Behaviors with a lower level of evidence for their association with opioid misuse include [48; 54; 55]:

- Aggressive demands for more drug
- Asking for specific medications
- Stockpiling medications during times when pain is less severe
- Using pain medications to treat other symptoms
- Reluctance to decrease opioid dosing once stable
- In the earlier stages of treatment:
  - Increasing medication dosing without provider permission
  - Obtaining prescriptions from sources other than the pain provider
  - Sharing or borrowing similar medications from friends/family

## INTERVENTIONS FOR SUSPECTED OR KNOWN ADDICTION OR DRUG DIVERSION

There are a number of actions that prescribers and dispensers can take to prevent or intervene in cases of drug diversion. These actions can be generally categorized based on the various mechanisms of drug diversion.

Prevention is the best approach to addressing drug diversion. As noted, the most common source of nonmedical use of prescribed opioids is from a family member or friend, through sharing, buying, or stealing. To avoid drug sharing among patients, healthcare professionals should educate patients on the dangers of sharing opioids and stress that “doing prescription drugs” is the same as “using street drugs” [49]. In addition, patients should be aware of the many options available to treat chronic pain aside from opioids. To prevent theft, patients should be advised to keep medications in a private place and to refrain from telling others about the medications being used.

Communication among providers and pharmacies can help to avoid inappropriate attainment of prescription drugs through “doctor shopping.” Prescribers should keep complete and up-to-date records for all controlled substance prescribing. When possible, electronic medical records should be integrated between pharmacies, hospitals, and managed care organizations [49]. If available, it is also best practice to periodically request a report from the state’s prescription reporting program to evaluate the prescribing of opioids to your patients by other providers [49].

When dealing with patients suspected of drug seeking/diversion, first inquire about prescription, over-the-counter, and illicit drug use and perform a thorough examination [49; 56]. Pill counting and/or UDT may be necessary to investigate possible drug misuse. Photo identification or other form of identification and social security number may be required prior to dispensing the drug, with proof of identity documented fully. If a patient is displaying suspicious behaviors, consider prescribing for limited quantities [56].

If a patient is found to be abusing prescribed opioids, this is considered a violation of the treatment agreement and the clinician must make the decision whether or not to continue the therapeutic relationship. If the relationship is terminated, it must be done ethically and legally. The most significant issue is the risk of patient abandonment, which is defined as ending a relationship with a patient without consideration of continuity of care and without providing notice to the patient. The American Medical Association Code of Ethics states that physicians have an obligation to support continuity of care for their patients. While physicians have the option of withdrawing from a case, they should notify the patient (or authorized decision maker) long enough in advance to permit the patient to secure another physician and facilitate transfer of care when appropriate [57]. Patients may also be given resources and/or recommendations to help them locate a new clinician.

Patients with chronic pain found to have an ongoing substance abuse problem or addiction should be referred to a pain specialist for continued treatment. Theft or loss of controlled substances is reported to the DEA. If drug diversion has occurred, the activity should be documented and a report to law enforcement should be made [58].

---

## COMPLIANCE WITH STATE AND FEDERAL LAWS

---

In response to the rising incidence in prescription opioid abuse, addiction, diversion, and overdose since the late 1990s, the FDA has mandated opioid-specific REMS to reduce the potential negative patient and societal effects of prescribed opioids. Other elements of opioid risk mitigation include FDA partnering with other governmental agencies, state professional licensing boards, and societies of healthcare professionals to help improve prescriber knowledge of appropriate and safe opioid prescribing and safe home storage and disposal of unused medication [24].

Several regulations and programs at the state level have been enacted in an effort to reduce prescription opioid abuse, diversion, and overdose, including [59]:

- Physical examination required prior to prescribing
- Tamper-resistant prescription forms
- Pain clinic regulatory oversight
- Prescription limits
- Prohibition from obtaining controlled substance prescriptions from multiple providers
- Patient identification required before dispensing
- Immunity from prosecution or mitigation at sentencing for individuals seeking assistance during an overdose

## CONTROLLED SUBSTANCES LAWS/RULES

The U.S. Drug Enforcement Administration (DEA) is responsible for formulating federal standards for the handling of controlled substances. In 2011, the DEA began requiring every state to implement electronic databases that track prescribing habits, referred to as PDMPs. Specific policies regarding controlled substances are administered at the state level [60].

According to the DEA, drugs, substances, and certain chemicals used to make drugs are classified into five distinct categories or schedules depending upon the drug's acceptable medical use and the drug's abuse or dependency potential [61]. The abuse rate is a determinate factor in the scheduling of the drug; for example, Schedule I drugs are considered the most dangerous class of drugs with a high potential for abuse and potentially severe psychologic and/or physical dependence.

## STATE-SPECIFIC LAWS AND RULES

Most states have established laws and rules governing the prescribing and dispensing of opioid analgesics. It is each prescriber's responsibility to have knowledge of and adhere to the laws and rules of the state in which he or she prescribes.

---

## MANAGEMENT OF OPIOID USE DISORDER

---

Management of opioid dependence entails different methods to achieve different goals, depending on the health situation and treatment history of the patient. These treatment approaches include [62]:

- Crisis intervention: Directed at immediate survival by reversing the potentially lethal effects of overdose with an opioid antagonist.
- Harm reduction: Intended to reduce morbidity and mortality associated with use of dirty needles and overdose.

- Detoxification/withdrawal: Aims to remove the opioid of abuse from the patient's body, either through gradual taper and substitution of a long-acting opioid or through ultra-rapid opioid detoxification.
- Maintenance treatment or opioid (agonist) replacement therapy: Aimed at reduction/elimination of illicit opioid use and lifestyle stabilization. Maintenance follows detoxification/withdrawal, whereby the patient is tapered from short-acting opioids and introduced to a long-acting opioid agonist, such as methadone or buprenorphine. Patients remain on agonist therapy short-term, long-term, or indefinitely depending on individual needs.
- Abstinence-oriented therapy: Treatment directed at cure. The patient is tapered off of short-acting opioids during the detoxification/withdrawal process and may be placed on an opioid antagonist with the goal of minimizing relapse.

All treatment approaches share the common goal of improving health outcomes and reducing drug-related criminality and public nuisance [62].

## CRISIS INTERVENTION

In response to acute overdose, the short-acting opioid antagonist naloxone is considered the criterion standard. Naloxone is effective in reversing respiratory depression and coma in patients who have overdosed. There is no evidence that subcutaneous or intramuscular use is inferior to intravenous naloxone. This prompted discussion of making naloxone available to the general public for administration outside the healthcare setting to treat acute opioid overdose, and in 2014, the FDA approved naloxone as an autoinjector dosage form for home use by family members or caregivers [63]. The autoinjector delivers 0.4 mg naloxone intramuscularly or subcutaneously. The autoinjector comes with visual and voice instruction, including directs to seek emergency medical care after use [63]. In 2015, the FDA approved intranasal naloxone after a fast-track designation and

priority review. Intranasal naloxone is indicated for the emergency treatment of known or suspected opioid overdose, as manifested by respiratory and/or central nervous system depression. It is available in a ready-to-use 2-mg, 4-mg, or 8-mg single-dose sprayer [64; 65; 87].

## HARM REDUCTION

Harm reduction measures are primarily employed to minimize the morbidity and mortality from opioid abuse and to reduce public nuisance [20; 66]. As a part of this effort, measures to prevent and minimize the frequency and severity of overdoses have been identified. Enrollment in opioid substitution therapy, with agents such as methadone and buprenorphine, substantially reduces the risk of overdose as well as the risk for infection and other sequelae of illicit opioid use [20; 66].

## DETOXIFICATION AND WITHDRAWAL

The process of tapering patients with opioid dependence from agonist therapy is often referred to as detoxification, or more accurately, medically supervised withdrawal [67; 68]. Its purpose is to eliminate physical dependence on opioid medications. It can be considered the medically supported transition to a medication-free state or to antagonist therapy. A careful and thorough review of the risks and benefits of detoxification should be provided, and informed consent obtained from patients prior to choosing this option [68; 69]. Detoxification alone should not be considered a treatment and should only be promoted in the context of a well-planned relapse-prevention program [62; 68].

Discontinuation of opioid use must be implemented slowly and cautiously to avoid a marked abstinence syndrome. Withdrawal symptoms may not begin for days after abrupt discontinuation of methadone or buprenorphine given their longer half-lives. Protracted abstinence, or post-acute withdrawal, may last for several months and is characterized by asthenia, depression, and hypotension. Post-acute withdrawal is more likely to occur with methadone than other opioids [67].

The three primary treatment modalities used for detoxification are opioid agonists, non-opioid medications, and rapid and ultra-rapid opioid detoxification [67]. The most frequently employed method of opioid withdrawal is a slow, supervised detoxification during which an opioid agonist, usually methadone, is substituted for the abused opioid [70]. Methadone is the most frequently used opioid agonist due to the convenience of its once-a-day dosing [67]. Methadone is highly bound to plasma proteins and accumulates more readily than heroin in all body tissues. Methadone also has a longer half-life, approximately 22 hours, which makes withdrawal more difficult than from heroin. Substitution therapy with methadone has a high initial dropout rate (30% to 90%) and an early relapse rate. Alternative pharmacologic detoxification choices include clonidine (with or without methadone), midazolam, trazodone, or buprenorphine [70].

Many opioid withdrawal symptoms, such as restlessness, rhinorrhea, lacrimation, diaphoresis, myosis, piloerection, and cardiovascular changes, are mediated through increased sympathetic activation, the result of increased neuron activity in the locus coeruleus. Non-opioid agents (such as clonidine), which inhibit hyperactivation of noradrenergic pathways stemming from the locus coeruleus nucleus, have been used to manage acute withdrawal [70; 71]. The first non-opioid treatment approved for the management of opioid withdrawal symptoms is lofexidine [86]. In studies, lofexidine resulted in less severe withdrawal symptoms and greater treatment retention than placebo.

However, some withdrawal symptoms, including anxiety and myalgias, are resistant to clonidine; benzodiazepines and nonsteroidal anti-inflammatory agents may be necessary to treat these symptoms. To mitigate withdrawal symptoms and assist in detoxification, alpha<sub>2</sub>-agonists, opioid agonist-antagonists, benzodiazepines, and antidepressants have been used [70].

Following detoxification, patients may feel exhausted and weak. Other complications, such as slight variations in hemodynamic status and gastrointestinal tract symptoms, follow quickly and may take several days to resolve. Muscle cramps and low back pain can be treated with nonsteroidal anti-inflammatory drugs. However, the newer cyclooxygenase-2 (COX-2) inhibitors may be advantageous because they produce fewer gastrointestinal side effects [70]. Insomnia is a frequent aspect of acute and protracted withdrawal, as opioids disrupt the normal sleep-wake cycle and many addicts require narcotics to sleep. Although long-term disruption of the normal sleep-wake cycle cannot be corrected rapidly, melatonin (3 mg), benzodiazepines, or antihistamines can be used with beneficial effects. Hypnosis and relaxation techniques are nonpharmacologic methods that may also be used [70]. Psychosocial treatments offered in addition to pharmacologic detoxification treatments positively impact treatment retention and completion, results at follow-up, and compliance [72; 73].

### **AGONIST REPLACEMENT OR ABSTINENCE THERAPY**

Two principle treatment modalities are offered for patients with opioid dependence: agonist maintenance or detoxification followed by outpatient or residential drug-free treatment. Both can be effective, with no clear indication for each, although agonist maintenance leads to greater treatment retention [74]. A reasonable approach is initial outpatient or residential treatment referral for patients relatively new to treatment, with agonist maintenance appropriate for patients with history of treatment failures, greater disease severity, or a history of drug overdoses. Naltrexone is best reserved for patients with strong legal incentives to abstain, family involvement to monitor treatment, or concurrent enrollment and involvement in a psychosocial intervention [75].

At present, there are no direct interventions that are capable of reversing the effects of drugs of dependence on learning and motivation systems [76]. Instead, the management of opioid dependence often consists of pharmacotherapy with methadone and buprenorphine, which do not eliminate physical dependence on opioids. These medications instead reduce the use of illicit opioids and produce very strong positive health outcomes as measured by decreased mortality, improved mental and physical health, and reduced risk of disease transmission [76]. Considering the high rate of relapse after detoxification, maintenance therapy with methadone or buprenorphine is currently considered to be the first-line treatment for patients with opioid dependence [62].

Any treatment for opioid dependence must take into consideration the chronic relapsing nature of opioid dependence, characterized by a variable course of relapse and remission in many patients. Treatments should emphasize patient motivation, psychoeducation, continuity of care, integration of pharmacotherapy and psychosocial support, and improved liaison between the treatment staff and the judicial system. Pharmacotherapy must be offered in a comprehensive healthcare context that also addresses the psychosocial aspects of dependence [62]. Patients who are dependent on opioids frequently suffer from physical and psychiatric disorders, and targeted interventions of psychiatric comorbidity are essential in improving treatment outcome for these patients [62]. Poly-substance abuse is the rule rather than the exception in opioid dependence, and concurrent use of other substances should be carefully monitored and treated when necessary [62]. Incarceration should never automatically result in discontinuation of an existing treatment; imprisonment offers a window of opportunity to initiate or restart treatment with a necessary continuation after release [62].

## Agonist Replacement Therapy

The goal of opioid replacement therapy is to reduce illicit drug use and associated health risks, with secondary goals of reducing unsafe sexual practices, improving vocational and psychosocial functioning, and enhancing quality of life [67]. The theoretical basis of opioid replacement stems from the finding that chronic opioid use results in an endogenous opioid deficiency as a result of the down-regulation of opioid production. This creates overwhelming cravings and necessitates interventions that shift the patient's attention and drive from obsessive preoccupation with the next use of opioids to more adaptive areas of focus, such as work, relationships, and non-drug leisure activities [67].

The neurobiologic changes resulting from prolonged opioid exposure provide a rationale for specific pharmacotherapies, such as long-acting opioid agonists, that are aimed at stabilizing these complex systems [77]. Opioid agonist maintenance treatment stabilizes brain neurochemistry by replacing short-acting opioids, which can create rapid changes in opioid levels in the serum and brain, with a long-acting opioid that has relative steady-state pharmacokinetics. Opioid agonist maintenance treatment is designed to have minimal euphoric effect, block the euphoria associated with administration of exogenous opioids (competitive antagonism), eliminate the risk of infectious disease and health consequences associated with injection drug use, and prevent opioid withdrawal [77].

Successful maintenance treatment entails stabilization of opioid dependence through opioid receptor occupation. Positron emission tomography studies have revealed that only 25% to 35% of brain opioid receptors are occupied during steady-state methadone maintenance, suggesting that unoccupied opioid receptors disrupted during cycles of opioid abuse could normalize during methadone main-

tenance [67]. Additionally, opioid replacement therapy blocks much of the euphoria from illicit heroin use. Long-term opioid agonist treatment also has a positive impact on public health, through significantly reducing overdose deaths, criminal activity, and the spread of infectious disease [67].

As of 2019, there were 1,691 treatment programs including opioid replacement therapy in the United States [78]. However, this represents only an estimated 19% of all patients with opioid use disorder. Although some have criticized the practice of methadone and buprenorphine therapy on the grounds that one opioid is merely being substituted for another, the clinical benefits strongly support this treatment modality [67]. When compared to active street heroin users, these benefits include a four-times lower HIV seroprevalence rates, 70% fewer crime-days per year, and a one-year mortality rate of 1% (versus 8%) [79].

## Abstinence-Oriented Therapies

The primary goal of abstinence-oriented interventions is cure, which is defined as long-term, stable abstinence from all opioids. Abstinence is achieved in two phases: detoxification and relapse prevention. Outcomes in abstinence-oriented programs are generally poor [62].

The primary goal of pharmacotherapy during detoxification is to alleviate opioid withdrawal severity and associated distress/medical complications and to enhance patient motivation to continue treatment. Withdrawal can also be reduced by psychosocial measures, such as contingency management or counseling, and as discussed, the addition of psychosocial therapy to pharmacologic treatment increases efficacy. Buprenorphine and clonidine are both used to manage withdrawal symptoms, but buprenorphine's advantages, compared with clonidine, are related to its favorable side effect profile and positive effects on well-being and psychosocial variables [62].

## 12-Step/Self-Help Programs

Twelve-step programs for opioid abuse and dependence include Narcotics Anonymous (NA), Heroin Anonymous (HA), and Methadone Anonymous (MA) and are modeled after Alcoholics Anonymous (AA), an abstinence-based support and self-improvement program that is based on the 12-step model of recovery. AA has helped hundreds of thousands of alcoholics achieve sobriety [80]. The 12-step model emphasizes acceptance of dependence as a chronic, progressive disease that can be arrested through abstinence but not cured. Additional elements include spiritual growth, personal responsibility, and helping other addicted persons. By inducing a shift in the consciousness of the addict, 12-step programs offer a holistic solution and are a resource for emotional support [80]. Although research on efficacy and patient outcomes in NA and MA is very limited, many prominent researchers emphasize the important role ongoing involvement in 12-step programs plays in recovery from substance abuse [81].

The understanding of drug dependence as a chronic and relapsing disorder has helped professionals gain a better comprehension of the vital role played by 12-step programs. Every patient attempting to recover from a substance use disorder will encounter a time when he or she faces urges to use without the resources or assistance of healthcare professionals. Twelve-step programs are not considered treatment, nor are they intended as substitutes for treatment. Instead, they are organizations that provide ongoing and indefinite support in the achievement and maintenance of abstinence and in personal growth and character development [81].

Part of the effectiveness of NA, HA, and MA is related to their ability to provide a competing and alternative reinforcer to drug use. Involvement in 12-step programs can enhance the quality of social support and the social network of the member, a potentially highly reinforcing aspect the person stands to forfeit if they resume drug using.

Other reinforcing elements of 12-step involvement include recognition for increasingly durable periods of abstinence and frequent awareness of the consequences of drug and alcohol use through attendance of meetings [82]. Research shows that establishing a pattern of 12-step program attendance early in treatment predicts the level of ongoing involvement. Emphasis and facilitation of early engagement in a 12-step program involvement are key [83].

## STIGMA OF ADDICTION

Many terms used in discussions of opioid use and misuse may have ambiguous meanings, and the absence of consensus in the terminology and definitions of substance use, substance use disorders, and addiction has led to considerable confusion and misconceptions. These misconceptions may be harbored by clinicians, patients, family members, and the public and can negatively impact patient interaction, assessment, treatment, and outcomes. This, coupled with pervasive stereotypes about what an opioid addict "looks" like, can negatively impact willingness to receive treatment or seek help and impair the patient's self-worth and mental health. Correction of these erroneous beliefs and attitudes is important, as is the use of nonpejorative and nonstigmatizing language when describing opioid analgesics, the patients who need them, and patients who develop aberrant behaviors or addiction involving opioids [32; 42]. It is important for all healthcare professionals to remember that addiction can affect any patients, regardless of age, sex, socioeconomic status, education, ability, or race.

## PROGNOSIS OF TREATMENT FOR OPIOID USE DISORDER

The relapse rate among patients receiving treatment for opioid dependence and other substance abuse is high (25% to 97%), comparable to that of other patients with chronic relapsing conditions, including hypertension and asthma [84]. Many cases of relapse are attributable to treatment non-compliance and lack of lifestyle modification [85].

Duration of agonist replacement therapy is usually recommended as a minimum of one year, and some patients will receive agonist replacement therapy indefinitely. Longer durations of treatment are associated with higher rates of abstinence from illicit opioids [76].

Much remains unknown about patient outcomes following termination of long-term opioid replacement therapy. Some patients aim to achieve total abstinence from all opioids, but little is known about patient characteristics and strategies used among those who remain abstinent. It is likely that at least some of the patients who remain abstinent from all opioids do so with the help of a 12-step support program, such as NA [76].

---

## CONCLUSION

---

Opioid analgesic medications can bring substantial relief to patients suffering from pain. However, the inappropriate use, abuse, and diversion of prescription drugs in America, particularly prescription opioids, has increased dramatically in recent years and has been identified as a national public health epidemic. A set of clinical tools, guidelines, and recommendations are now available for prescribers who treat patients with opioids. By implementing these tools, the clinician can effectively address issues related to the clinical management of opioid prescribing, opioid risk management, regulations surrounding the prescribing of opioids, and problematic opioid use by patients. In doing so, healthcare professionals are more likely to achieve a balance between the benefits and risks of opioid prescribing, optimize patient attainment of therapeutic goals, and avoid the risk to patient outcome, public health, and viability of their own practice imposed by deficits in knowledge.

**Works Cited**

1. Federation of State Medical Boards. *Guidelines for the Chronic Use of Opioid Analgesics*. Washington, DC: The Federation of State Medical Boards; 2017.
2. Institute of Medicine, Committee on Advancing Pain Research, Care, and Education. *Relieving Pain in America: A Blueprint for Transforming Prevention, Care, Education, and Research*. Washington, DC: The National Academies Press; 2011.
3. Kenen K, Mack K, Paulozzi L. Trends in prescriptions for oxycodone and other commonly used opioids in the United States, 2000–2010. *Open Medicine*. 2012;6(2):e41.
4. Owen GT, Burton AW, Schade CM, Passik S. Urine drug testing: current recommendations and best practices. *Pain Physician*. 2012;15:ES119-ES133.
5. Groenewald CB, Rabbits JA, Gebert JT, Palermo TM. Trends in opioid prescriptions among children and adolescents in the United States: a nationally representative study from 1996 to 2012. *Pain*. 2016;157(5):1021-1027.
6. Miller NS, Greenfeld A. Patient characteristics and risks factors for development of dependence on hydrocodone and oxycodone. *Am J Ther*. 2004;11:26-32.
7. Substance Abuse and Mental Health Services Administration. *Drug Abuse Warning Network, 2006: National Estimates of Drug-Related Emergency Department Visits*. Rockville, MD: U.S. Department of Health and Human Services; 2007.
8. Substance Abuse and Mental Health Services Administration. *Drug Abuse Warning Network, 2011: National Estimates of Drug-Related Emergency Department Visits*. Rockville, MD: Substance Abuse and Mental Health Services Administration; 2014.
9. Management of Opioid Therapy for Chronic Pain Working Group. *VA/DoD Clinical Practice Guideline for Opioid Therapy for Chronic Pain*. Washington, DC: Department of Veterans Affairs, Department of Defense; 2017.
10. Dowell D, Haegerich TM, Chou R. CDC guideline for prescribing opioids for chronic pain—United States, 2016. *MMWR*. 2016;65(1):1-49.
11. Chou R, Fanciullo GJ, Fine PG, et al. Clinical guidelines for the use of chronic opioid therapy in chronic noncancer pain. *J Pain*. 2009;10(2):113-130.
12. Centers for Disease Control and Prevention. About CDC's Opioid Prescribing Guideline. Available at <https://www.cdc.gov/drugoverdose/prescribing/guideline.html>. Last accessed April 20, 2021.
13. Argoff CE, Silvershein DI. A comparison of long- and short-acting opioids for the treatment of chronic noncancer pain: tailoring therapy to meet patient needs. *Mayo Clin Proc*. 2009;84(7):602-612.
14. McCarberg BH, Barkin RL. Long-acting opioids for chronic pain: pharmacotherapeutic opportunities to enhance compliance, quality of life, and analgesia. *Am J Ther*. 2001;8(3):181-186.
15. National Comprehensive Cancer Network. Adult Cancer Pain, 2021. Available at [https://www.nccn.org/professionals/physician\\_gls/pdf/pain.pdf](https://www.nccn.org/professionals/physician_gls/pdf/pain.pdf). Last accessed April 20, 2021.
16. U.S. Food and Drug Administration. Medication Guides: Distribution Requirements and Inclusion in Risk Evaluation and Mitigation Strategies (REMS). Available at <https://www.fda.gov/media/79776/download>. Last accessed April 20, 2021.
17. Manchikanti L, Abdi S, Atluri S, et al. American Society of Interventional Pain Physicians (ASIPP) guidelines for responsible opioid prescribing in chronic non-cancer pain: part 2—guidance. *Pain Physician*. 2012;15(3 Suppl):S67-S116.
18. Cheattle MD. Risk Assessment: Safe Opioid Prescribing Tools. Available at <https://www.practicalpainmanagement.com/resource-centers/opioid-prescribing-monitoring/risk-assessment-safe-opioid-prescribing-tools>. Last accessed April 21, 2021.
19. Butler SF, Budman SH, Fernandez KC, Fanciullo GJ, Jamison RN. Cross-validation of a Screener to Predict Opioid Misuse in Chronic Pain Patients (SOAPP-R). *J Addict Med*. 2009;3(2):66-73.
20. Singer JA. Harm Reduction: Shifting from a War on Drugs to a War on Drug-Related Deaths. Available at <https://www.cato.org/policy-analysis/harm-reduction-shifting-war-drugs-war-drug-related-deaths>. Last accessed April 25, 2021.
21. Fine PG, Finnegan T, Portenoy RK. Protect your patients, protect your practice: practical risk assessment in the structuring of opioid therapy in chronic pain. *J Fam Pract*. 2010;59(9 Suppl 2):S1-S16.
22. Katz NP. Opioid Prescribing Toolkit: A Workbook for Clinicians. New York, NY: Oxford University Press; 2010.
23. American Chronic Pain Association. ACPA – Stanford Resource Guide to Pain Management: 2020 Edition. Available at <https://www.theacpa.org/wp-content/uploads/2020/03/ACPA-Resource-Guide-2020-2-26-2020.pdf>. Last accessed April 22, 2021.
24. Strickland JM, Huskey A, Brushwood DB. Pharmacist-physician collaboration in pain management practice. *J Opioid Manag*. 2007;3:295-301.
25. Brushwood DB. Screening Controlled Substance Prescriptions for Legitimacy: The VIGIL System. Available at [http://www.empaa.org/downloads/EMPAAP2010/presentation/BrushwoodDavid\\_VIGIL\\_EMPAA2010.pdf](http://www.empaa.org/downloads/EMPAAP2010/presentation/BrushwoodDavid_VIGIL_EMPAA2010.pdf). Last accessed April 22, 2021.
26. Miner J, Babitz M, Dunn A, Fondario A, Smith M. Utah Clinical Guidelines on Prescribing Opioids for Treatment of Pain. Salt Lake City, UT: Utah Department of Health; 2018.
27. American Pain Foundation. Breakthrough cancer pain: mending the break in the continuum of care. *J Pain and Palliat Care Pharmacother*. 2011;25(3):252-264.

## #55151 Responsible and Effective Opioid Prescribing

---

28. Kaye AD, Jones MR, Kaye AM, et al. Prescription opioid abuse in chronic pain: an updated review of opioid abuse predictors and strategies to curb opioid abuse (part 2). *Pain Physician*. 2017;20(Supp 2):S111-S133.
29. American Society of Anesthesiologists. Opioid Abuse. Available at <https://www.asahq.org/madeforthismoment/pain-management/opioid-treatment/opioid-abuse>. Last accessed April 22, 2021.
30. National Comprehensive Cancer Network. NCCN Guidelines: Treatment by Cancer Type. Available at [https://www.nccn.org/guidelines/category\\_1](https://www.nccn.org/guidelines/category_1). Last accessed April 21, 2021.
31. Federal Drug Administration. Safe Opioid Disposal: Remove the Risk Outreach Toolkit. Available at <https://www.fda.gov/drugs-ensuring-safe-use-medicine/safe-opioid-disposal-remove-risk-outreach-toolkit>. Last accessed April 25, 2021.
32. Oliver J, Coggins C, Compton P, et al. American Society for Pain Management Nursing position statement: pain management in patients with substance use disorders. *Pain Manage Nurs*. 2012;13(3):169-183.
33. U.S. Food and Drug Administration. FDA Analysis of Long-Term Trends in Prescription Opioid Analgesic Products: Quantity, Sales, and Price Trends. Available at <https://www.fda.gov/media/111695/download>. Last accessed April 25, 2021.
34. Dalal S, Bruera E. Assessment and management of pain in the terminally ill. *Prim Care Clin Office Pract*. 2011;38:195-223.
35. Gao W, Gulliford M, Higginson IJ. Prescription patterns of analgesics in the last 3 months of life: a retrospective analysis of 10,202 lung cancer patients. *Br J Cancer*. 2011;104(11):1704-1710.
36. Hudgins JD, Porter JJ, Monuteaux MC, Bourgeois FT. Trends in opioid prescribing for adolescents and young adults in ambulatory care settings. *Pediatrics*. 2019;143(6):e20181578.
37. Breuer B, Fleishman SB, Cruciani RA, Portenoy RK. Medical oncologists' attitudes and practice in cancer pain management: a national survey. *J Clin Oncol*. 2011;29(36):4769-4775.
38. Abraham JL. *A Physician's Guide to Pain and Symptom Management in Cancer Patients*. 3rd ed. Baltimore, MD: Johns Hopkins University Press; 2014.
39. Lovegrove MC, Dowell D, Geller AI, et al. US emergency department visits for acute harms from prescription opioid use, 2016–2017. *Am J Pub Health*. 2019;109:784-791.
40. Centers for Disease Control and Prevention. CDC Advises Against Misapplication of the Guideline for Prescribing Opioids for Chronic Pain. Available at <https://www.cdc.gov/media/releases/2019/s0424-advises-misapplication-guideline-prescribing-opioids.html>. Last accessed April 20, 2021.
41. Loeser JD, Schatzman ME. Chronic pain management in medical education: a disastrous omission. *Postgrad Med*. 2017;129(3):332-335.
42. New York State Department of Health. Implantable Infusion Pumps for Non-Cancer Pain. Available at [https://www.health.ny.gov/health\\_care/medicaid/redesign/2016/medtronic\\_sources\\_dla\\_rev.htm](https://www.health.ny.gov/health_care/medicaid/redesign/2016/medtronic_sources_dla_rev.htm). Last accessed April 20, 2021.
43. Mailis A, Taenzer P. Evidence-based guideline for neuropathic pain interventional treatments: spinal cord stimulation, intravenous infusions, epidural injections and nerve blocks. *Pain Res Manage*. 2012;17(3):150-158.
44. National Institute on Drug Abuse. Benzodiazepines and Opioids. Available at <https://www.drugabuse.gov/drugs-abuse/opioids/benzodiazepines-opioids>. Last accessed April 24, 2021.
45. Pergolizzi JV Jr, Mercadante S, Echaburu AV, et al. The role of transdermal buprenorphine in the treatment of cancer pain: an expert panel consensus. *Curr Med Res Opin*. 2009;25(6):1517-1528.
46. Passik SD, Kirsh KL, Whitcomb L, et al. A new tool to assess and document pain outcomes in chronic pain patients receiving opioid therapy. *Clin Ther*. 2004;26:552-561.
47. Atluri SL, Akbik H, Sudarshan G. Prevention of opioid abuse in chronic non-cancer pain: an algorithmic, evidence-based approach. *Pain Physician*. 2012;15:ES177-ES189.
48. National Institute on Drug Abuse. Minimizing the Misuse of Prescription Opioids in Patients with Chronic Nonmalignant Pain. Available at <https://www.drugabuse.gov/nidamed/centers-excellence/resources/minimizing-misuse-prescription-opioids-in-patients-chronic-nonmalignant-pain>. Last accessed April 24, 2021.
49. American College of Preventive Medicine. Use, Abuse, Misuse and Disposal of Prescription Pain Medication Patient Guide. Available at <http://www.yuma.usmc-mccs.org/mccsyuma/assets/File/Drug%20Facts%20Sheets/painmedsclinicalreference.pdf>. Last accessed April 25, 2021.
50. U.S. Food and Drug Administration. Disposal of Unused Medicines: What You Should Know. Available at <https://www.fda.gov/drugs-safe-disposal-medicines/disposal-unused-medicines-what-you-should-know>. Last accessed April 25, 2021.
51. Substance Abuse and Mental Health Services Administration. Results from the 2019 National Survey on Drug Use and Health: Detailed Tables. Available at <https://www.samhsa.gov/data/report/2019-nsduh-detailed-tables>. Last accessed April 24, 2021.
52. Sekhon R, Aminjavahery N, Davis CN Jr, et al. Compliance with opioid treatment guidelines for chronic non-cancer pain (CNCP) in primary care at a Veterans Affairs Medical Center (VAMC). *Pain Med*. 2013;14(10):1548-1556.
53. Meier B. Increase in Urine Testing Raises Ethical Questions. Available at [https://www.nytimes.com/2013/08/02/business/increase-in-urine-testing-raises-ethical-questions.html?\\_r=0](https://www.nytimes.com/2013/08/02/business/increase-in-urine-testing-raises-ethical-questions.html?_r=0). Last accessed April 24, 2021.
54. Passik SD. Issues in long-term opioid therapy: unmet needs, risks, and solutions. *Mayo Clin Proc*. 2009;84:593-601.

55. Holliday S, Hayes C, Dunlop A. Opioid use in chronic non-cancer pain. Part 2: prescribing issues and alternatives. *Australian Family Physician*. 2013;42:104-111.
56. U.S. Department of Justice Office of Diversion Control. Don't Be Scammed By A Drug Abuser. Available at <https://www.deadiversion.usdoj.gov/pubs/brochures/drugabuser.htm>. Last accessed April 24, 2021.
57. American Medical Association. Code of Ethics: Opinion 1.1.5 Terminating the Patient-Physician Relationship. Available at <https://www.ama-assn.org/sites/ama-assn.org/files/corp/media-browser/code-of-medical-ethics-chapter-1.pdf>. Last accessed April 24, 2021.
58. U.S. Department of Health and Human Services Drug Diversion: What Is a Prescriber's Role in Preventing the Diversion of Prescription Drugs? Available at <https://www.hhs.gov/guidance/document/drug-diversion-what-prescribers-role-preventing-diversion-prescription-drugs>. Last accessed April 24, 2021.
59. Centers for Disease Control and Prevention. Risk for overdose from methadone used for pain relief—United States, 1999–2010. *MMWR*. 2012;61(26):493-497.
60. Hannon K. The Prescription Drug Crisis in New York State: A Comprehensive Approach. Available at <https://www.scribd.com/doc/82474334/Prescription-Drug-Abuse-Crisis-in-NYS-Comprehensive-Approach-New>. Last accessed April 24, 2021.
61. U.S. Drug Enforcement Administration. Drug Scheduling. Available at <https://www.dea.gov/drug-scheduling>. Last accessed April 24, 2021.
62. van den Brink W, Haasen C. Evidence-based treatment of opioid-dependent patients. *Can J Psychiatry*. 2006;51(10):635-646.
63. Cassels C. FDA Okays Handheld Autoinjector for Opioid Overdose. Available at <https://www.medscape.com/viewarticle/823039>. Last accessed April 24, 2021.
64. Narcan Nasal Spray (naloxone intranasal) [package insert]. Available at [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2015/208411lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2015/208411lbl.pdf). Last accessed April 24, 2021.
65. U.S. Food and Drug Administration. FDA Moves Quickly to Approve Easy-To-Use Nasal Spray to Treat Opioid Overdose. Available at <https://wayback.archive-it.org/7993/20180125101447/https://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm473505.htm>. Last accessed April 24, 2021.
66. Darke S, Hall W. Heroin overdose: research and evidence-based intervention. *J Urban Health*. 2003;80(2):189-200.
67. Krantz MJ, Mehler PS. Treating opioid dependence: growing implications for primary care. *Arch Intern Med*. 2004;164(3):277-288.
68. Substance Abuse and Mental Health Services Administration. *Federal Guidelines for Opioid Treatment Programs*. HSS Publication No. (SMA) PEP15-FEDGUIDEOTP. Rockville, MD: Substance Abuse and Mental Health Services Administration; 2015.
69. Substance Abuse and Mental Health Services Administration. SAMHSA Opioid Overdose Prevention Toolkit. HHS Publication No. (SMA) 16-4742. Rockville, MD: Substance Abuse and Mental Health Services Administration; 2016.
70. Kaye AD, Gevirtz C, Bosscher HA, et al. Ultrarapid opiate detoxification: a review. *Can J Anaesth*. 2003;50(7):663-671.
71. American Society of Addiction Medicine. National practice guideline for the use of medications in the treatment of addiction involving opioid use. Available at <https://www.asam.org/docs/default-source/practice-support/guidelines-and-consensus-docs/asam-national-practice-guideline-supplement.pdf>. Last accessed April 25, 2021.
72. Amato L, Minozzi S, Davoli M, Vecchi S, Ferri M, Mayet S. Psychosocial and pharmacological treatments versus pharmacological treatments for opioid detoxification. *Cochrane Database Syst Rev*. 2008;(4):CD005031.
73. Amato L, Minozzi S, Davoli M, Vecchi S. Psychosocial and pharmacological treatments versus pharmacological treatments for opioid detoxification. *Cochrane Database Syst Rev*. 2011;9:CD005031.
74. Fiellin DA, O'Connor PG. Office-based treatment of opioid-dependent patients. *N Engl J Med*. 2002;347(11):817-823.
75. Nunes EV, Sullivan MA, Levin FR. Treatment of depression in patients with opiate dependence. *Biol Psychiatry*. 2004;56(1):793-802.
76. Wasan AD, Correll DJ, Kissin I, O'Shea S, Jamison RN. Iatrogenic addiction in patients treated for acute or subacute pain: a systematic review. *J Opioid Manag*. 2006;2(1):16-22.
77. Fiellin DA, Friedland GH, Gourevitch MN. Opioid dependence: rationale for and efficacy of existing and new treatments. *Clin Infect Dis*. 2006;43(Suppl 4):S173-S177.
78. Substance Abuse and Mental Health Services Administration. *National Survey of Substance Abuse Treatment Services (N-SSATS): 2019. Data on Substance Abuse Treatment Facilities*. Rockville, MD: Substance Abuse and Mental Health Services Administration; 2020.
79. Kleber HD. Methadone maintenance 4 decades later: thousands of lives saved but still controversial. *JAMA*. 2008;300(19):2303-2305.
80. Humphreys K, Wing S, McCarty D, et al. Self-help organizations for alcohol and drug problems: toward evidence-based practice and policy. *J Substance Abuse Treat*. 2004;26(3):151-165.
81. Ronel N, Gueta K, Abramsohn Y, Caspi N, Adelson M. Can a 12-step program work in methadone maintenance treatment? *Int J Offender Ther Comp Criminol*. 2011;55(7):1135-1153.

## #55151 Responsible and Effective Opioid Prescribing

---

82. Stoops WW, Lile JA, Glaser PE, Hays LR, Rush CR. Alternative reinforcer response cost impacts cocaine choice in humans. *Prog Neuropsychopharmacol Biol Psychiatry*. 2012;36(1):189-193.
83. Donovan DM, Ingalsbe MH, Benbow J, Daley DC. 12-step interventions and mutual support programs for substance use disorders: an overview. *Soc Work Public Health*. 2013;28(3-4):313-332.
84. Dixon DW. Opioid Abuse Follow-Up. Prognosis. Available at <https://emedicine.medscape.com/article/287790-followup#e4>. Last accessed April 25, 2021.
85. Vukmir RB. Drug seeking behavior. *Am J Drug Alcohol Abuse*. 2004;30:551-575.
86. U.S. Food and Drug Administration. FDA Approves the First Non-Opioid Treatment for Management of Opioid Withdrawal Symptoms in Adults. Available at <https://www.fda.gov/news-events/press-announcements/fda-approves-first-non-opioid-treatment-management-opioid-withdrawal-symptoms-adults>. Last accessed April 25, 2021.
87. U.S. Food and Drug Administration. FDA Approves Higher Dosage of Naloxone Nasal Spray to Treat Opioid Overdose. Available at <https://www.fda.gov/news-events/press-announcements/fda-approves-higher-dosage-naloxone-nasal-spray-treat-opioid-overdose>. Last accessed October 11, 2021.

### Evidence-Based Practice Recommendations Citation

Manchikanti L, Abdi S, Atluri S, et al. American Society of Interventional Pain Physicians (ASIPP) guidelines for responsible opioid prescribing in chronic non-cancer pain: part 2—guidance. *Pain Physician*. 2012;15(3 Suppl):S67-S116. Available at <https://painphysicianjournal.com/2012/july/2012;%2015;S67-S116.pdf>. Last accessed April 28, 2021.